

# Field Evaluation of Almond Varieties

---

## Project No.: 17-Hort2-Lampinen

**Project Leaders:** Bruce Lampinen, Dept. of Plant Sciences, University of California, One Shields Ave., Davis, CA 95616 (email: [bdlampinen@ucdavis.edu](mailto:bdlampinen@ucdavis.edu)) and Phoebe Gordon (UCCE Madera and Merced Counties) Roger Duncan (UCCE Stanislaus County), Luke Milliron (UCCE Butte Glenn Tehama Counties)

**Project Cooperators:** Tom Gradziel, Sam Metcalf, Maria Contador, Sabrina Marchand, Misha Marchand (UCD), Joe Connell (UCCE Butte County), Chico State University, Salida School District, and Creekside Farming Company

### Summary:

The next generation Regional Almond Variety Trials were planted in the winter of 2014 in Butte, Stanislaus and Madera counties. Rows of Nonpareil were alternated with 29 varieties and/or selections at all 3 sites. Trees at the Butte, Stanislaus and Madera trial were planted on Krymsk 86, Nemaguard and Hansen 536 rootstocks, respectively, (with the exceptions listed at the bottom of Table 5). Unlike the previous generation Regional Almond Variety Trials, there are four replications of each of the varieties and selections at each of the three sites in the 2014 trials. Bloom overlap of pollenizers with Nonpareil was generally good at all the sites with the exception of UCD 3-40. Yields in 2020 were higher than in past years, primarily due to excellent weather during bloom. Main kernel defects observed in 2020 were doubles, twins, naval orange worm damage, blanks and severe shrivel.

**Objective:** The objective is to evaluate new almond varieties and selections in replicated trials at three locations in the almond growing areas of California.

### Results and discussion:

#### General observations for each site

#### Butte.

The average February rainfall in Chico is 4.4 inches but in 2020, no rainfall was measured at the nearest (Durham) CIMIS station. Following excellent bloom density and weather, yields were very high in Chico despite observations of mediocre bee activity. UCD 18-20 which showed some leafing delay and failure symptoms in 2018, displayed these symptoms again in 2020. No bloom or spring foliar diseases were observed in 2020. In the summer of 2018, almond leaf scorch (ALS) was confirmed on Self Fruitful P16.013 and UCD 1-271 by Dr. Lindsey Burbank at USDA-ARS in Parlier. In 2019 ALS symptoms were observed on Self-fruitful P16.013 and Booth. In 2020, ALS symptoms, which in some areas were severe, were observed on all varieties/selections and observed across the field. Because of harvest sequence, irrigation was kept off in the trial for over 1 month (from well before the first shaking of varieties on August 10, until the final pick up on September 11). Because of this extreme drought there was significant defoliation both from water stress and mites by September 11. Mummy counts were highest (averaging

200 nuts per tree or more) in Y117-86-03, Winters, and UCD 7-159, the later two of which never reached 100% hull split before being shaken – likely because of hull-tights induced by the extreme water stress. The extensive early defoliation led to the pushing of new leaves across the orchard in October. Trunk/scaffold cankers (especially band canker) are most prevalent in UCD 7-159, Wood Colony, and Sterling, followed by Folsom, and Nonpareil. The heavy crop and original poor scaffold selection (all scaffolds originating from the same plane) resulted in tree loss in several varieties, most notably in Aldrich. Kester on Hansen and Eddie have had extensive tree loss because of gophers, as well as unknown causes.

### Stanislaus.

Bloom weather in 2020 was dry and mild with no rain from start to finish, leading to the best yields in this trial so far. There were no obvious signs of bloom or spring foliar diseases in this trial. Severe hull rot (*Rhizopus*) on Y121-42-99, Folsom, Kester on Hansen rootstock (but not Nemaguard), and 8-201. Moderate hull rot (mostly *Rhizopus*) on 8-160, Nonpareil, 1-232, 8-27, and Eddie. *Aspergillus niger* hull rot on 1-232. Overall, growth of trees were better than in past years, although there were widespread signs of moderate leaf chlorosis / mottling presumably from alkaline irrigation water and use of Nemaguard rootstock. Trees on Hansen rootstock were much better. Foliar symptoms of moderate zinc and potassium deficiency were evident in some areas/ varieties which were confirmed by leaf tissue analyses. Leaf samples indicated elevated levels of chloride in the trial.

### Madera

Bloom conditions were excellent in 2020, with little to no observed diseases. Due to the extremely high nut set, there was some bending and breaking of upright primary and secondary scaffolds. Several scattered trees have died due to gopher damage that is most likely from previous years, and some of the stressed trees also suffered from shothole borer infestations before succumbing. Hull rot continues to be an issue (Fig. 1), and canopy closure in addition to hull rot is resulting in the loss of some lower limbs on all varieties. Some leaf-footed bug and stink bug were observed in the orchard at the end of 2020, and these pests will be monitored for in 2021.

## **Bloom, Hullsplit, Yield and Quality**

### **2020**

Butte- UCD 3-40 bloomed considerably earlier than anything else but it is being removed from data collection since it is too early to serve as a pollinizer in these trials. Bloom was fairly compact at the Butte site in 2020 with only about 6 days of difference between full bloom dates for the earliest versus latest varieties (Fig. 2). Bloom overlap was generally good across all varieties and selections. Hullsplit ranged from July 12<sup>th</sup> to September 1<sup>st</sup> in 2020 (Fig. 3). Midday canopy PAR interception ranged from 50 to 83% with Nonpareil coming in at 74% (Table 2). Yield ranged from 1741 kernel pounds per acre for Kester on Hansen to 4659 for Nonpareil (Table 3). Yield per unit PAR intercepted ranged from 28.6 for Kester on Hansen to 63.3 for Aldrich (Table 4). Cumulative yield for the Butte site from 2017-2020 ranged from 4896 for UCD1-271 to 13035 for Nonpareil (Table 5).

Stanislaus- Bloom was also fairly compact with good bloom overlap at the Stanislaus site with full bloom ranging from February 17 to February 22. Hullsplit ranged from July 12 to August 31<sup>st</sup> at the Stanislaus trial in 2020 (Fig. 3). Midday PAR interception varied from 38.3% for UCD8-160 to 66.9% for Kester on Hansen rootstock (Table 6). Yields ranged from 1453 kernel pounds per acre for Y121-43-99 to 3726 for Kester on Hansen rootstock (Table 7). Yield per unit PAR intercepted was very high for some varieties at this site in 2020 ranging from 27.8 for UCD8-27 to 78.7 for Winters (Table 8). Cumulative yield ranged from 5237 kernel pounds per acre for UCD8-27 to 10,828 for Kester on Hansen (Table 9).

Madera- As at the other two sites, bloom was compact at the Madera site in 2020 with full bloom ranging 8 days from Feb. 17 to Feb. 25 (UCD 3-40 was earlier but it is no longer being monitored in these trials). Hullsplit ranged from July 7 to September 6 (Fig. 2). Midday PAR interception ranged from 56% for UCD8-160 to 89% for Folsom in 2020 (Table 10). Eight varieties had PAR interception greater than 80% which is the maximum we recommend so shading is becoming an issue in this trial. Yields in 2020 ranged from 1799 kernel pounds per acre for UCD1-271 to 5004 kernel pounds per acre for Nonpareil (Table 11). Yield per unit PAR intercepted ranged from 22.1 for UCD1-271 to 61.1 kernel pounds per 1% PAR intercepted for Nonpareil (Table 12). Cumulative yields ranged from 4045 for UCD3-40 to 13,446 kernel pounds per acre for Nonpareil (Table 13). Leaf tissue analysis for the Madera site in 2020 are shown in Table 16.

Average cumulative yield for all three sites averaged ranged from 5465 for UCD1-271 to 11,667 for Nonpareil (Table 14). UCD18-20 which is the second top yielding selection or variety overall also has a large number of doubles every year so this may be problematic. Although yields in all 3 trials were significantly higher in 2020, the values are in the same range as our previous McFarland trial in Kern County and significantly higher than in the previous generation trials (Fig. 4).

#### **Outreach activities:**

In January 2021, Luke Milliron gave the talk “Almond Variety Evaluation in the Sacramento Valley” at the UCCE Sacramento Valley Almond Grower Meeting.

#### **Materials and methods:**

##### Regional Almond Variety Trials Planted in 2014

The next generation almond variety trials were planted in the winter of 2014 in Butte (Chico State University), Stanislaus (Salida School District Site), and Madera (Chowchilla grower site) counties. The varieties and selections planted are listed in Table 1. The first 30 items are common to all 3 sites and a few different items added at individual sites are listed at the bottom of Table 1. Trees at the Butte, Stanislaus and Madera trial were planted on Krymsk 86, Nemaguard and Hansen 536 rootstocks respectively (with the exceptions listed at the bottom of Table 1). Trees were planted at a spacing of 18' x 22' at the Butte site (110 trees/acre), 16' x 21' at the Stanislaus site (130 trees/acre) and 12' x 21' at the Madera site (173 trees/acre). These densities are significantly higher than the previous generation RAVTs where planting densities for the Butte, San Joaquin and Kern trials were 64, 75 and 86 trees per acre respectively. Of

the items planted in the main trials, fourteen are either partially or fully self-fertile (Table 1).

Bloom, hullsplit, canopy light interception and yield data collection were initiated in 2016. Bloom data were collected approximately every three days and recorded as onset of bloom, full bloom, and the end of petalfall. Hullsplit was recorded from the beginning of the first non-blank splits to completion of hullsplit.

**Publications that emerged from this work:**

Gordon, P.; Duncan, R.; Milliron, L.; Lampinen, B. (2020). Field Evaluation of Almond Varieties: A Look at Regional Trial Results through Sixth Leaf. *West Coast Nut*. September 17. <http://www.wcngg.com/2020/09/17/field-evaluation-of-almond-varieties/>

Gradziel, T.; Milliron, L. (2020). Breeding pt. 3: Almond with Tom Gradziel. *Growing the Valley*. February 18. <https://www.growingthevalleypodcast.com/podcastfeed/almond>

Acknowledgements

The authors wish to thank the Almond Board of California for their continued support of this project. In addition, we would like to thank Chico State University/Jeff Boles, the Salida School District/Lane Parker as well as the Creekside Farming Company for their cooperation on the next generation variety trials.

Table 1. Varieties and selections planted at the next generation regional almond variety trials. Items 1-29 are planted at all 3 sites while additional material planted at individual sites is listed at the end. Trees at the Butte, Stanislaus and Madera sites were planted on Krymsk 86, Nemaguard and Hansen 536 rootstock respectively (exceptions are noted at bottom of table).



#	Variety or selection	Self-fertile*	Source
1	Eddie		Bright's
2	Capitola		Burchell
3	Supareil		Burchell
4	Self-fr P13-019***	yes	Burchell
5	Self-fr P16-013***	yes	Burchell
6	Booth		Burchell
7	Sterling		Burchell
8	Bennett		Duarte
9	Nonpareil		Fowler
10	Durango		Fowler
11	Jenette		Fowler
12	Aldrich		Fowler
13	Winters	partial	UCD
14	Sweetheart	partial	UCD
15	Kester (2-19E)*		UCD
16	UCD3-40***		UCD
17	UCD18-20		UCD
18	UCD1-16		UCD
19	UCD8-160	yes	UCD
20	UCD8-27	yes	UCD
21	UCD1-271	yes	UCD
22	UCD1-232	yes	UCD
23	UCD7-159	yes	UCD
24	UCD8-201	yes	UCD
25	Y121-42-99	yes	USDA
26	Y117-86-03	yes	USDA
27	Yorizane (Y116-161-99)**	yes	USDA
28	Y117-91-03	yes	USDA
29	Folsom		Wilson
30	Wood Colony on Krymsk 86 (Butte site only)		
31	Wood Colony on Nemaguard (Madera site only planted one year later after Lone Star was removed)		

\*Kester was planted at all three sites on the usual rootstock for the site. In addition, at the Butte and Stanislaus sites it was also planted in the replicated trial on Hansen 536 rootstock.

\*\*Y116-161-99 was released as Yorizane in 2020

\*\*\*Self-fruitful P16.013 and Self-fruitful P13.019 were eliminated from data collection in 2020 since they have been dropped by the nursery that developed them.

Table 2. 2020 canopy PAR interception for Butte County.

	#reps	Variety or selection	PAR interception	
			(%)	
Butte	4	Supareil	82.7	a
	4	Capitola	78.3	a b
	4	Sweetheart	75.0	a b c
	4	Folsom	74.6	a b c d
	4	UCD18-20	74.6	a b c d
	4	Kester	74.5	a b c d
	4	Nonpareil	74.1	a b c d
	4	Booth	73.8	a b c d
	4	Winters	73.6	a b c d
	4	Y117-91-03	73.3	a b c d e
	4	UCD3-40	72.6	a b c d e
	4	UCD1-16	72.1	a b c d e
	4	Durango	71.3	a b c d e f
	4	UCD8-27	71.2	a b c d e f
	4	Sterling	68.9	b c d e f g
	4	Y117-86-03	68.9	b c d e f g
	4	Aldrich	68.8	b c d e f g
	4	Bennett	68.0	b c d e f g h
	4	Eddie	63.9	c d e f g h
	4	UCD8-201	62.8	c d e f g h
4	UCD1-232	62.6	c d e f g h	
4	UCD7-159	61.9	d e f g h	
4	Jenette	61.6	d e f g h i	
4	Kester/Hansen	60.7	e f g h i	
4	Yorizane	59.2	f g h i	
4	UCD1-271	56.7	g h i	
4	Wood Colony	55.7	h i	
4	UCD8-160	49.7	i	

Table 3. 2020 yield for Butte County.

	#reps	Variety or selection	2020 Yield	
			(kernel lbs/ac)	
Butte	4	Nonpareil	4573	a
	4	Aldrich	4353	a b
	4	Capitola	4116	a b c
	4	Durango	3756	b c d
	4	Winters	3756	b c d
	4	UCD18-20	3746	b c d
	4	Booth	3576	c d e
	4	UCD7-159	3495	c d e f
	4	Y117-91-03	3423	c d e f
	4	Jenette	3367	c d e f g
	4	UCD1-16	3336	d e f g h
	4	Sweetheart	3300	d e f g h
	4	Wood Colony	3300	d e f g h
	4	Bennett	3233	d e f g h
	4	Yorizane	3229	d e f g h
	4	Sterling	3156	d e f g h i
	4	Supareil	3154	d e f g h i
	4	UCD8-201	3046	d e f g h i j
	4	Kester	2998	d e f g h i j
	4	UCD1-232	2900	e f g h i j
	4	Folsom	2898	e f g h i j
	4	Y117-86-03	2753	f g h i j
	4	Eddie	2595	g h i j
	4	UCD8-160	2567	g h i j
	4	UCD3-40	2543	h i j
	4	UCD1-271	2415	i j k
4	UCD8-27	2358	j k	
4	Kester/Hansen	1735	k	

Table 4. 2020 yield per unit light intercepted for Butte County.

	#reps	Variety or selection	Yield per unit PAR	
			intercepted	
Butte	4	Aldrich	63.3	a
	4	Nonpareil	61.7	a b
	4	Wood Colony	59.0	a b c
	4	UCD7-159	56.6	a b c d
	4	Jenette	54.8	b c d e
	4	Yorizane	54.6	b c d e f
	4	Durango	52.5	c d e f g
	4	Capitola	52.4	c d e f g
	4	UCD8-160	51.7	c d e f g h
	4	Winters	50.9	c d e f g h
	4	UCD18-20	50.2	c d e f g h
	4	UCD8-201	48.5	c d e f g h i
	4	Booth	48.5	d e f g h i
	4	Bennett	47.7	d e f g h i j
	4	Y117-91-03	46.8	d e f g h i j k
	4	UCD1-16	46.3	e f g h i j k
	4	UCD1-232	46.3	e f g h i j k
	4	Sterling	45.6	f g h i j k
	4	Sweetheart	44.0	g h i j k
	4	UCD1-271	42.8	h i j k l
	4	Kester	40.4	i j k l m
	4	Y117-86-03	40.2	i j k l m
	4	Eddie	39.6	i j k l m
	4	Folsom	39.1	j k l m
4	Supareil	38.1	k l m	
4	UCD3-40	35.1	l m n	
4	UCD8-27	33.2	m n	
4	Kester/Hansen	28.4	n	



Table 5. Cumulative yield for Butte County from 2017-2020.

	#reps	Variety or selection	Cumulative yield (kernel lbs/ac)	
<b>Butte</b>	4	Nonpareil	12949	a
	4	UCD18-20	11412	a b
	4	Booth	11312	b
	4	Aldrich	10989	b
	4	Jenette	10222	b c
	4	Y117-91-03	10103	b c d
	4	Durango	9944	b c d
	4	Winters	9923	b c d
	4	Capitola	9727	b c d e
	4	Yorizane	9061	c d e f
	4	UCD8-201	8979	c d e f
	4	UCD8-160	8694	c d e f g
	4	Folsom	8693	c d e f g
	4	Kester	8660	c d e f g
	4	Wood Colony	8654	c d e f g
	4	Bennett	8660	c d e f g
	4	Y117-86-03	8256	d e f g
	4	UCD1-232	8181	d e f g
	4	UCD1-16	8171	d e f g
	4	UCD7-159	7960	e f g
	4	Eddie	7908	e f g
	4	Sterling	7888	e f g
	4	UCD8-27	7438	f g
	4	Sweetheart	7429	f g
	4	Supareil	6964	g
	4	Kester/Hansen	6953	g
	4	UCD3-40	6940	g
	4	UCD1-271	4887	h

Table 6. PAR interception for Stanislaus site 2020.

	#reps	Variety or selection	PAR interception	
			(%)	
<b>Stanislaus</b>	4	Kester/Hansen	66.9	a
	4	Sweetheart	64.4	a b
	4	Supareil	63.3	a b c
	4	Y117-91-03	60.8	a b c d
	4	Booth	58.0	a b c d e
	4	Eddie	57.4	a b c d e f
	4	UCD3-40	55.5	a b c d e f g
	4	Capitola	54.6	b c d e f g
	4	Sterling	54.2	b c d e f g h
	4	UCD8-27	53.4	b c d e f g h
	4	Kester	51.2	b c d e f g h i
	4	UCD18-20	51.1	b c d e f g h i
	4	Folsom	50.5	c d e f g h i
	4	Bennett	50.3	d e f g h i
	4	Nonpareil	49.2	d e f g h i
	4	Jenette	48.7	d e f g h i
	4	UCD1-271	48.0	d e f g h i
	4	UCD8-201	47.0	d e f g h i
	4	Aldrich	46.5	e f g h i
	4	Durango	46.2	e f g h i
	4	UCD1-16	45.0	f g h i
	4	UCD1-232	44.1	f g h i
	4	Y121-42-99	43.9	f g h i
	4	UCD7-159	43.8	g h i
	4	Yorizane	42.5	g h i
	4	Y117-86-03	41.9	g h i
	4	Winters	40.3	h i
	4	UCD8-160	38.3	i

Table 7. 2020 yield for the Stanislaus site.

	#reps	Variety or selection	2020 Yield	
			(kernel lbs/ac)	
<b>Stanislaus</b>	4	Kester/Hansen	3703	a
	4	Nonpareil	3521	a b
	4	Aldrich	3098	a b c
	4	Capitola	3036	a b c d
	4	Y117-91-03	3009	a b c d
	3	Bennett	2978	a b c d
	4	Durango	2879	b c d e
	4	Eddie	2869	b c d e
	4	Supareil	2732	c d e f
	4	Booth	2701	c d e f
	4	Winters	2671	c d e f
	4	UCD7-159	2646	c d e f g
	4	UCD18-20	2568	c d e f g
	4	Y117-86-03	2531	c d e f g
	4	Sweetheart	2525	c d e f g
	4	Kester	2375	c d e f g
	4	Yorizane	2357	c d e f g
	4	Sterling	2350	c d e f g
	4	Folsom	2273	d e f g
	4	UCD1-16	2268	d e f g
	4	UCD1-232	2108	e f g h
	4	UCD8-160	2074	f g h
	3	UCD8-201	2064	f g h
	4	UCD3-40	2012	f g h
	4	UCD1-271	1975	f g h
	4	Jenette	1889	g h
	4	UCD8-27	1403	h i
	4	Y121-42-99	1089	i

Table 8. Yield per unit PAR intercepted for Stanislaus site 2020.

		Yield per unit PAR	
#reps	Variety or selection	intercepted	
4	Winters	78.1	a
4	Nonpareil	77.0	a
4	Aldrich	66.9	a b
4	Durango	62.3	a b c
4	UCD7-159	61.1	a b c d
4	Y117-86-03	60.6	a b c d e
3	Bennett	57.3	a b c d e f
4	Capitola	56.0	b c d e f
4	Kester/Hansen	55.8	b c d e f
4	Yorizane	55.0	b c d e f
4	UCD8-160	54.0	b c d e f
4	Folsom	50.8	b c d e f g
4	UCD18-20	50.5	b c d e f g
4	UCD1-16	50.2	b c d e f g
4	Eddie	50.1	b c d e f g
4	Y117-91-03	49.5	b c d e f g
4	UCD1-232	47.3	b c d e f g h
4	Booth	46.7	b c d e f g h
4	Kester	46.6	b c d e f g h
3	UCD8-201	43.8	c d e f g h
4	Sterling	43.5	c d e f g h
4	Supareil	42.9	c d e f g h
4	UCD1-271	41.3	c d e f g h
4	Sweetheart	39.2	d e f g h
4	Jenette	38.5	e f g h
4	UCD3-40	36.4	f g h
4	Y121-42-99	29.5	g h
4	UCD8-27	26.2	h

Stanislaus

Table 9. Cumulative yield for Stanislaus County from 2016-2020.

	#reps	Variety or selection	Cumulative yield	
			(kernel lbs/ac)	
<b>Stanislaus</b>	3	Kester/Hansen	11089	a
	3	Y117-91-03	9412	b
	4	UCD18-20	9290	b
	3	Bennett	8950	b c
	4	Nonpareil	8520	b c d
	4	UCD8-160	8353	b c d e
	4	Aldrich	8162	b c d e f
	4	UCD7-159	8129	b c d e f
	4	Booth	8103	b c d e f
	4	Capitola	8069	b c d e f
	4	Kester	7993	b c d e f
	3	Durango	7969	b c d e f
	4	Yorizane	7965	b c d e f
	4	Winters	7887	b c d e f
	4	Y117-86-03	7778	b c d e f g
	3	Sterling	7490	b c d e f g
	4	Eddie	7255	c d e f g
	3	UCD8-201	7167	c d e f g
	4	UCD1-232	6881	d e f g h
	4	Sweetheart	6806	d e f g h
	4	Folsom	6684	d e f g h
	4	Supareil	6644	d e f g h
	4	UCD1-271	6537	d e f g h
	4	UCD1-16	6496	e f g h
	4	Y121-42-99	6208	e f g h
	4	Jenette	6185	f g h
	4	UCD3-40	5867	g h
	4	UCD8-27	5151	h

Table 10. PAR interception for 2020 season for Madera site.

	#reps	Variety or selection	PAR interception	
			(%)	
<b>Madera</b>	4	Folsom	89.2	a
	4	Booth	85.9	a b
	4	Supareil	84.8	a b c
	4	Sterling	83.8	a b c d
	4	Eddie	83.2	a b c d e
	4	Capitola	82.2	a b c d e f
	4	Nonpareil	81.8	a b c d e f
	4	UCD1-271	81.8	a b c d e f
	4	Kester	79.9	a b c d e f g
	4	Sweetheart	77.2	a b c d e f g
	4	Aldrich	75.6	a b c d e f g h
	4	UCD3-40	75.4	a b c d e f g h
	4	Durango	74.1	a b c d e f g h
	4	UCD18-20	69.6	a b c d e f g h
	4	Bennett	69.5	a b c d e f g h
	4	Y117-86-03	68.8	a b c d e f g h
	4	Yorizane	67.7	b c d e f g h
	4	UCD8-27	67.7	b c d e f g h
	4	UCD7-159	65.9	c d e f g h
	4	Y117-91-03	63.8	c d e f g h
4	Jenette	63.5	c d e f g h	
4	Wood Colony	63.2	d e f g h	
4	UCD1-16	62.6	d e f g h	
4	Winters	61.8	e f g h	
4	UCD1-232	60.7	f g h	
4	UCD8-201	59.8	g h	
4	UCD8-160	55.6	h	

Table 11. 2020 yield for Madera site in 2020.

	#reps	Variety or selection	2020 Yield	
			(kernel lbs/ac)	
<b>Madera</b>	4	Nonpareil	5004	a
	4	Durango	3535	b
	4	Sterling	3470	b c
	4	Booth	3468	b c
	4	Supareil	3443	b c d
	4	Capitola	3337	b c d e
	4	Aldrich	3171	b c d e f
	4	Jenette	3022	b c d e f g
	4	Y117-86-03	3014	b c d e f g
	4	UCD7-159	2931	b c d e f g h
	4	Yorizane	2839	b c d e f g h i
	4	Kester	2809	b c d e f g h i
	4	UCD8-201	2806	b c d e f g h i
	4	Bennett	2787	c d e f g h i
	4	Eddie	2741	c d e f g h i
	4	Wood Colony	2721	d e f g h i
	4	UCD18-20	2695	e f g h i
	3	Sweetheart	2640	e f g h i
	4	Winters	2618	e f g h i
	4	UCD1-16	2555	f g h i
4	Folsom	2552	f g h i	
4	UCD8-160	2418	f g h i j	
4	Y117-91-03	2327	g h i j	
4	UCD8-27	2314	g h i j	
4	UCD3-40	2184	h i j	
4	UCD1-232	2143	i j	
4	UCD1-271	1799	j	

Table 12. Yield per unit PAR intercepted for Madera site in 2020.

	#reps	Variety or selection	Yield per unit PAR	
			intercepted	
<b>Madera</b>	4	Nonpareil	61.1	a
	4	UCD8-201	50.2	a b
	4	Durango	47.5	b c
	4	Jenette	47.2	b c
	4	Wood Colony	46.9	b c
	4	UCD7-159	45.0	b c d
	4	Yorizane	43.8	b c d
	4	Y117-86-03	43.2	b c d
	4	UCD8-160	41.6	b c d
	4	UCD1-16	40.9	b c d e
	4	Winters	40.3	b c d e
	4	Capitola	40.1	b c d e f
	4	Sterling	39.6	b c d e f
	4	Aldrich	39.5	b c d e f
	4	Booth	39.5	b c d e f
	4	Bennett	39.1	b c d e f
	4	Supareil	38.1	b c d e f
	4	UCD18-20	36.9	b c d e f
	4	Y117-91-03	36.6	c d e f
	4	Kester	34.7	c d e f
4	UCD1-232	34.7	c d e f	
4	UCD8-27	34.2	c d e f	
3	Sweetheart	34.0	c d e f	
4	Eddie	32.4	d e f g	
4	UCD3-40	27.8	e f g	
4	Folsom	27.0	f g	
4	UCD1-271	21.0	g	



Table 13. Cumulative yield for 2016-2020 for Madera site.

	#reps	Variety or selection	Cumulative yield	
			(kernel lbs/ac)	
<b>Madera</b>	4	Nonpareil	13446	a
	3	Yorizane	13021	a b
	4	Y117-86-03	12142	a b c
	4	UCD18-20	12118	a b c d
	4	Capitola	11307	a b c d e
	4	Kester	11260	a b c d e
	4	Booth	11176	a b c d e
	4	Jenette	11078	a b c d e
	4	Y117-91-03	10764	b c d e f
	3	Sweetheart	10372	c d e f g
	4	Bennett	10324	c d e f g
	4	UCD8-201	10148	c d e f g
	4	Eddie	10102	c d e f g
	4	Sterling	10061	c d e f g
	4	Aldrich	9855	c d e f g
	4	Winters	9777	c d e f g
	4	Durango	9699	c d e f g
	4	UCD1-16	9650	c d e f g h
	4	UCD8-160	9416	c d e f g h
	4	Folsom	9368	d e f g h
	4	Supareil	9292	e f g h
	4	UCD8-27	8349	f g h
	4	UCD7-159	7756	g h i
	3	UCD1-232	7034	h i
	4	Wood Colony	5374	i j
	3	UCD1-271	4836	i j
3	UCD3-40	3940	j	

Table 15. Main  
are

**Cumulative yield  
(lbs/acre)**

kernel defects for 2020 harvest. Items  
listed if they had 6% or more of kernels

Variety or selection	Cumulative yield (lbs/acre)	kernel defects for 2020 harvest. Items listed if they had 6% or more of kernels
Nonpareil	11638	a
UCD18-20	10940	a b exhibiting the defect.
Booth	10197	a b c
Y117-91-03	10140	a b c d
Yorizane	9742	a b c d e
Capitola	9701	a b c d e
Aldrich	9668	a b c d e
Kester/Hansen	9647	a b c d e f
Y117-86-03	9392	b c d e f
Bennett	9331	b c d e f
Durango	9316	b c d e f
Winters	9195	b c d e f
Jenette	9161	b c d e f
UCD8-201	8910	b c d e f g
UCD8-160	8821	c d e f g
Sterling	8570	c d e f g
Eddie	8422	c d e f g
Kester**	8374	c d e f g
Folsom	8245	c d e f g h
UCD1-16	8106	c d e f g h
Sweetheart	8005	d e f g h
Wood Colony*	7985	d e f g h
UCD7-159	7966	e f g h
Supareil	7723	e f g h
UCD1-232	7396	f g h i
UCD8-27	7049	g h i
Y121-42-99*	6208	i
UCD3-40	5731	i
UCD1-271	5473	i

\*Stanislaus site only

\*\*Butte and Madera sites

Varieties with defect	Trial					
	Butte	(%)	Stanislaus	(%)	Madera	(%)
Double kernels	UCD 8-27	51	UCD 8-27	24	UCD 8-27	29
	UCD 8-201	44	UCD 8-201	18	UCD 8-201	23
	UCD 1-16	17	UCD 1-16	12	Y117-86-03	18
	Folsom	16			Booth	16
	UCD18-20	15			UCD 1-16	11
	Booth	15			Folsom	8
	Y117-86-03	14			Capitola	7
	Y117-91-03	12			UCD18-20	7
	UCD1-232	9				
	UCD 8-160	8				
	Wood Colony	7				
	Nonpareil	7				
	Kester	6				
	Twin kernels (two kernels within the same pellicle)	UCD 8-27	14	UCD 8-27	15	UCD 8-27
Jenette		11	UCD 8-201	12	UCD1-232	10
UCD 8-201		9	Supareil	9.5	UCD 8-201	10
UCD 8-160		9	Sweetheart	9.5	Jenette	8
UCD1-232		6	UCD 3-40	9	Supareil	8
Booth		6	Booth	9	UCD 1-16	7
			UCD 8-160	7.5	UCD 3-40	6
			Nonpareil	7		
			Bennett	6		
			Folsom	6		
		Jenette	6			
Navel orange worm damage	UCD1-271	7			UCD1-271	12
	UCD 3-40	6			UCD 7-159	9
					UCD 8-27	7
					Y117-91-03	7
					UCD 3-40	7
					Bennett	6
					Eddie	6
					Winters	6
				Y117-86-03	6	
				Supareil	6	
Blank kernels	UCD1-232	10	UCD 8-27	7.5		
	UCD 8-27	6	Folsom	6.5		
	Y117-86-03	6				
Chipped/broken			UCD 8-27	7.5	UCD18-20	11
			Winters	7		
			Sterling	6.5		
			UCD 8-160	6		

Varieties with defect	Trial					
	Butte	(%)	Stanislaus	(%)	Madera	(%)
Crease	Jenette	17	Jenette	13		
	Capitola	14	UCD 7-159	10		
	Nonpareil	9	Folsom	8		
	Sterling	8	Capitola	7		
	Y117-86-03	7	Supareil	6.5		
	Folsom	6				
	UCD 7-159	6				
Shrivel	Capitola	14				
	Yorizane	9				
	Supareil	7				
	UCD 8-201	6				
Rupture/callous					UCD1-271	28
					Lonestar	14
					UCD1-232	14
					Eddie	14
					UCD 8-160	11
					Yorizane	10
					Y117-91-03	8
				Nonpareil	6	
Stain/discolor	UCD1-271	27	UCD 1-271	21	UCD1-232	24
	Yorizane	10	Yorizane	21	UCD1-271	24
	UCD 8-160	9	Bennett	6.5	Eddie	20
	Winters	7	Eddie	4.5	Sweetheart	12
	Kester/Hansen	6	UCD 1-232	4	UCD 8-160	12
					Nonpareil	11
					UCD 3-40	10
					Bennett	10
					Kester	9
					UCD 8-27	9
					Y117-86-03	9
					UCD 8-201	8
					Capitola	6
					Lonestar	6
				Yorizane	6	
mold	Eddie	16	UCD 1-232	28		
	UCD1-271	15	UCD 3-40	21		
	Bennett	10	Y117-86-03	21		
	UCD1-232	9	P16.013	20		
	Kester/Hansen	7	UCD 1-271	18		
	Nonpareil	7	Bennett	7.5		
	UCD 7-159	7	UCD 8-160	7		
	Folsom	6	Y117-91-03	6		
	Wood Colony	6				
gum	UCD 3-40	14			Capitola	13
					UCD 8-27	8

Table 16. Nutrient Levels in July-Sampled Leaves. Stanislaus County Regional Almond Variety Trial 2020

	N (%)	P (%)	K (%)	Ca (%)	Mg (%)
1-271	2.68 a	0.14 abc	1.94 abc	4.13 bcde	1.23 abc
Folsom	2.63 ab	0.13 abcde	1.65 bcde	3.85 cde	1.18 abcd
Kester/Hansen	2.58 abc	0.13 abcdef	1.34 e	5.10 a	1.21 abc
8-27	2.57 abc	0.14 a	1.73 abcde	3.77 cde	1.08 cdef
P13-019	2.54 abcd	0.13 abcdefg	1.57 bcde	4.19 bcd	1.18 cdef
Sterling	2.54 abcd	0.14 abcd	1.60 bcde	3.77 cde	1.03 def
Sweetheart	2.54 abcd	0.14 ab	1.80 abcde	2.97 f	0.92 f
Y116-161-99	2.48 abcde	0.12 bcdefg	2.01 ab	3.98 bcde	1.14 cde
Kester	2.48 abcde	0.12 efg	1.35 e	3.91 cde	1.13 cde
Y121-42-99	2.47 abcde	0.12 abcdefg	1.97 abc	3.48 ef	1.06 cdef
18-20	2.47 abcde	0.12 bcdefg	1.49 cde	3.64 def	1.13 cde
1-16	2.46 abcde	0.12 abcdefg	1.71 abcde	3.48 ef	0.99 ef
Winters	2.45 abcde	0.13 abcdef	1.36 e	4.27 bcd	1.34 a
8-160	2.45 abcde	0.12 abcdefg	1.54 bcde	4.46 abc	1.18 abcd
1-232	2.44 abcde	0.13 abcdef	1.41 de	4.08 bcde	1.19 abcd
P16.013	2.43 abcde	0.12 abcdefg	1.73 abcde	3.92 bcde	1.17 bcd
Booth	2.42 abcde	0.12 abcdefg	1.89 abcd	3.75 de	1.13 cde
Y117-91-03	2.39 bcde	0.13 abcdef	2.18 a	4.08 bcde	1.15 bcde
Nonpareil	2.38 bcde	0.11 fg	1.75 abcde	3.94 bcde	1.04 def
8-201	2.38 bcde	0.12 defg	1.79 abcde	3.83 cde	1.12 cde
Capitola	2.37 bcde	0.12 defg	1.61 bcde	4.17 bcde	1.31 ab
Eddie	2.35 cde	0.12 abcdefg	1.73 abcde	4.00 bcde	1.21 abc
3-40	2.35 cde	0.12 abcdefg	1.58 bcde	4.62 ab	1.20 abcd
Aldrich	2.34 cde	0.11 g	1.47 bcd	4.22 cde	1.11 cde
Bennett	2.33 cde	0.13 abcdefg	1.70 abcde	4.01 bcde	1.07 cdef
Durango	2.32 cde	0.11 efg	1.36 e	3.95 bcde	1.19 abcd
Supareil	2.29 de	0.12 abcdefg	1.66 bcde	3.84 cde	1.07 cdef
Y117-86-03	2.26 e	0.12 cdefg	1.62 bcde	3.78 cde	1.09 cde
7-159	2.26 e	0.12 abcdefg	1.79 abcde	4.21 bcd	1.17 abcd
Jennette	2.25 e	0.13 abcdefg	1.73 abcde	3.95 bcde	1.13 cde
<b>Critical value</b>	<b>2.2 - 2.5</b>	<b>0.1 – 0.3</b>	<b>&gt; 1.4</b>	<b>&gt; 2.0</b>	<b>&gt; 0.25</b>

Values followed by the same letters are statistically similar (Tukeys  $P \leq 0.05$ )

Table 16 (continued).

	Zn (ppm)	Mn (ppm)	Cl ( %)	Na (%)
1-271	18.6 ab	33.3 a	0.26 bcd	0.11 abc
Folsom	13.2 bcd	33.3 a	0.29 abcd	0.08 bc
Kester/Hansen	20.0 a	47.4 a	0.07 e	0.04 c
8-27	12.6 cd	40.6 a	0.31 abcd	0.13 abc
P13-019	14.6 abcd	38.7 a	0.33 abcd	0.10 abc
Sterling	15.8 abcd	34.9 a	0.19 de	0.07 bc
Sweetheart	15.3 abcd	28.2 a	0.23 d	0.06 c
Y116-161-99	14.9 abcd	33.0 a	0.31 abcd	0.16 ab
Kester	11.8 cd	38.4 a	0.32 abcd	0.12 abc
Y121-42-99	13.6 bcd	37.3 a	0.27 bcd	0.06 c
18-20	11.5 cd	27.6 a	0.27 bcd	0.18 a
1-16	10.1 d	31.9 a	0.24 cd	0.07 bc
Winters	14.4 bcd	33.6 a	0.30 abcd	0.17 a
8-160	10.9 cd	36.4 a	0.21 de	0.06 c
1-232	10.9 cd	43.0 a	0.27 bcd	0.08 bc
P16.013	12.9 cd	38.3 a	0.21 de	0.08 bc
Booth	14.9 abcd	31.7 a	0.29 abcd	0.13 abc
Y117-91-03	16.0 abc	27.4 a	0.33 abcd	0.14 ab
Nonpareil	12.4 cd	44.2 a	0.25 bcd	0.08 bc
8-201	10.9 cd	30.5 a	0.28 abcd	0.09 bc
Capitola	16.1 abc	37.0 a	0.32 abcd	0.11 abc
Eddie	12.2 cd	45.5 a	0.29 abcd	0.09 bc
3-40	14.4 bcd	48.4 a	0.26 bcd	0.15 ab
Aldrich	11.9 cd	25.1 a	0.42 a	0.18 a
Bennett	13.9 bcd	49.8 a	0.25 bcd	0.11 abc
Durango	13.1 cd	27.1 a	0.39 ab	0.10 abc
Supareil	11.5 cd	37.6 a	0.19 de	0.06 c
Y117-86-03	10.6 cd	29.8 a	0.33 abcd	0.19 a
7-159	11.4 cd	33.0 a	0.29 abcd	0.09 bc
Jennette	11.4 cd	33.4 a	0.38 abc	0.18 a
<b>Critical value</b>	<b>&gt;15</b>	<b>&gt;20</b>	<b>&lt;0.3</b>	<b>&lt;0.25</b>

## Leaf nutrient observation summary:

- There were statistically significant differences in all nutrients among varieties, although not always agronomically important.
- Aldrich tends to be lower than average in N, P, K, Zn and Mg but high in Cl and Na
- The three varieties with the highest potassium were USDA varieties Y117-91-03, Yorizane & Y121-42-99 (over 2%)
- Kester on Nemaguard and Hansen were numerically the lowest in K (1.3%)
- Most varieties were deficient in Zn (< 15ppm).
- Hansen increased zinc levels in Kester from 11.8 ppm to 20.0 ppm, compared to Nemaguard
- Chloride toxicity is an emerging problem in this trial. Aldrich, Durango and Jenette have the highest chloride levels while Supareil and Sterling had the lowest
- Kester on Nemaguard had 0.32 % chloride compared to 0.07% for Kester on Hansen rootstock

Table 17. Leaf tissue analyses for Madera RA VT in 2020.

	Nitrogen			P			Potassium	
Booth	2.60	a	Sweetheart	0.144	a	Folsom	3.03	a
Kester	2.51	ab	Folsom	0.143	ab	UCD1-271	3.00	ab
Folsom	2.49	abc	Sterling	0.142	abc	Capitola	2.87	abc
Sweetheart	2.48	abc	Booth	0.140	abc	Booth	2.77	abcd
Eddie	2.47	abcd	UCD3-40	0.137	abc	Y117-91-03	2.74	abcd
UCD1-271	2.41	abcde	UCD1-271	0.136	abc	Sterling	2.72	abcd
Nonpareil	2.38	abcdef	Kester	0.136	abc	Eddie	2.75	abcd
UCD8-27	2.37	abcdef	Eddie	0.136	abc	Jenette	2.68	abcd
Capitola	2.36	abcdef	Nonpareil	0.129	abc	Kester	2.66	abcd
UCD3-40	2.36	abcdef	Wood Colony	0.126	abc	Durango	2.65	abcd
UCD18-20	2.33	abcdef	UCD8-27	0.126	abc	Y117-86-03	2.60	abcd
Sterling	2.30	abcdef	Supareil	0.125	abc	Wood Colony	2.54	abcd
UCD1-16	2.28	bcdef	Y117-86-03	0.125	abc	Supareil	2.52	abcd
Aldrich	2.26	bcdef	UCD7-159	0.125	abc	Nonpareil	2.51	abcd
UCD1-232	2.26	bcdef	Aldrich	0.124	abc	UCD8-201	2.47	abcde
Jenette	2.24	bcdef	UCD1-16	0.124	abc	UCD7-159	2.43	abcde
Y117-86-03	2.23	bcdef	Jenette	0.124	abc	UCD1-232	2.41	abcde
Durango	2.23	bcdef	Capitola	0.123	abc	Aldrich	2.34	abcde
Wood Colony	2.22	bcdef	UCD1-232	0.122	abc	UCD18-20	2.29	abcde
Winters	2.20	cdef	UCD8-201	0.122	abc	Bennett	2.28	abcde
UCD8-201	2.20	cdef	Durango	0.120	abc	UCD8-27	2.22	bcde
UCD7-159	2.20	cdef	UCD18-20	0.119	abc	Yorizane	2.19	cde
Bennett	2.19	cdef	UCD8-160	0.119	abc	Sweetheart	2.17	cde
Yorizane	2.16	def	Bennett	0.117	bc	UCD8-160	2.12	cde
UCD8-160	2.16	def	Yorizane	0.116	c	UCD3-40	2.05	de
Supareil	2.15	ef	Y117-91-03	0.116	c	UCD1-16	2.01	de
Y117-91-03	2.07	f	Winters	0.116	c	Winters	1.69	e

Fig. 1. Number of hullrot strikes per tree at the Madera site in 2020.




Fig. 2. Bloom data for 2020 by site and variety or selection.






Fig. 3. Hullsplit by site, variety and selection for 2020.




Fig. 4. Average annual yield for all varieties and selections combined at each trial by orchard age. Kern, Butte old and Delta are from the previous generation variety trials and the McFarland trial was in Kern County with Mario Viveros.




Fig. 5. Google Earth images of the three sites. Note extensive tree loss in several areas at the Madera trial and Stanislaus trials.

