

TREE PEST UPDATES

75 Santa Barbara Rd., 2nd floor, Pleasant Hill, CA 94523-4488 (925) 646-6540
724 Third St., Brentwood, CA 94513 (925) 634-3012

April 26, 2002

PEACH TWIG BORER

HOST CROPS: Almonds, Apricots, Peaches, Nectarines, Plums

BIOFIX: Flight started in most orchards between **April 7th and 11th**. If you had traps in your orchard, use the date when you first started to catch moths as your own orchard biofix.

TREATMENT AND TIMING: In-season sprays may be needed to protect fruit and nuts from damage if you have significant trap counts. This will be especially important if you did not apply a dormant spray or bloom sprays (BT or Success). Effective control can be achieved by applying most sprays 400-500 degree days (DD) after biofix. This is projected to occur **May 10-20 OR 33-40 days after your own orchard biofix**. The table below lists some of the materials which can be used to control PTB in the various crops. BT and Success have very low human toxicity and are less likely to disrupt beneficial insects.

BT (*Bacillus thuringiensis*) is an organically acceptable option and will be more effective if applied at 300 DD and again between 450-500 DD. 300 DD is projected to occur **May 3 OR 26 days after your own orchard biofix**. 450-500 DD is projected to occur **May 13-17 OR 36-40 days after your own orchard biofix**.

Material	Almond	Apricot	Nectarine	Peach	Plum
Ambush/Pounce 1,2,3	X			X	
Asana ^{1,2,3}	X	X	X	X	X
BT	X	X	X	X	X
Diazinon ^{2,3}		X	X	X	X
Imidan ^{1,2,3}	X	X	X	X	X
Guthion ^{1,2,3}	X		X	X	X
Sevin ^{1,2,3}	X	X	X	X	X
Success	X	X	X	X	X
Supracide ^{1,2,3}	X	X	X	X	X

¹ toxic to mite predators - may increase mite problems

² toxic to general predators and parasites – may lead to secondary pest outbreaks

³ restricted material permit or applicator's certificate required for use

Mating Disruption materials are also available and effective in controlling PTB in orchards larger than 5-10 acres. They should have been applied before biofix and should be reapplied at the recommended interval for as long as you need your fruit protected. Think about this option for next year.

PEACH TWIG BORER UPDATE

UC Cooperative Extension
75 Santa Barbara Rd, 2nd floor
Pleasant Hill, CA 94523

Janet Caprile
Farm Advisor

UNITED STATES DEPARTMENT OF AGRICULTURE, UNIVERSITY OF CALIFORNIA AND ALAMEDA/CONTRA COSTA COUNTIES COOPERATING

The University of California prohibits discrimination against or harassment of any person on the basis of race, color, national origin, religion, sex, physical or mental disability, medical condition, (cancer-related or genetic characteristics), ancestry, marital status, age, sexual orientation, citizenship, or status as a covered veteran (special disabled veteran, Vietnam-era veteran or any other veteran who served on active duty during a war or in a campaign or expedition for which a campaign badge has been authorized). University policy is intended to be consistent with the provisions of applicable State and Federal Laws. Inquiries regarding the University's nondiscrimination policies may be directed to the Affirmative Action/Staff Personnel Services Director, university of California, Agriculture and Natural Resources, 300 Lakeside Drive, Oakland, CA 94612-3550 [Telephone: (510) 987-0096].

What's a Biofix?: It's just the beginning of the flight for each new generation. We usually have 3 generations for peach twig borer in this area. We use the Biofix to begin degree day calculations for each generation so we know when egg laying, hatchout, and other lifecycle events will happen. This helps us to time our treatments most effectively.

What's a Degree Day? Insects develop faster or slower depending on the temperature. Degree days are a measure based on the maximum and minimum temperatures for each day which allow us to figure out how fast the insects are developing. You may see them abbreviated as DD or °D. If you have the daily maximum & minimum temperatures for your orchard, you can look the degree days up on a chart. If you have access to the Internet, you can get Brentwood weather data and do a degree day calculation from the UC IPM Program home page. This page also lets you calculate the projected degree days based on historical weather data so you can make projections for potential treatment windows (this is how I do it!). The address is <http://www.ipm.ucdavis.edu>. Give me a call if you would like a degree day chart or more information

